

SEMIPROJECTIVITY OF UNIVERSAL C^* -ALGEBRAS GENERATED BY ALGEBRAIC ELEMENTS

TATIANA SHULMAN

ABSTRACT. Let p be a polynomial in one variable whose roots either all have multiplicity more than 1 or all have multiplicity exactly 1. It is shown that the universal C^* -algebra of a relation $p(x) = 0$, $\|x\| \leq 1$ is semiprojective. In the case of all roots multiple it is shown that the universal C^* -algebra is also residually finite-dimensional. Applications to polynomially compact operators are given.

INTRODUCTION

The "general lifting problem" in C^* -algebras is a class of problems which are formulated in the following way: for a quotient C^* -algebra A/I and its element b with some specific properties, to find (or to prove the existence of) a preimage of b in A with the same properties. Lifting problems are closely related to the theory of stability of relations which, roughly speaking, investigates when "an almost representation of a relation" is "close" to a representation (for strict definitions see [5]).

By a relation we mean a system of equalities of the form $f(x_1, \dots, x_n, x_1^*, \dots, x_n^*) = 0$, where f is a non-commutative polynomial (an element of the corresponding free algebra), which can include also inequalities of the form $\|x_i\| \leq c$ or $\|x_i\| < c$ (they also can be written as polynomial equalities if one extends the list of generators).

Recall that a C^* -algebra D is *projective* if for any C^* -algebra A , its ideal I and every $*$ -homomorphism $\phi : D \rightarrow A/I$, there exists $\tilde{\phi}$ such that the diagram

$$\begin{array}{ccc} & & A \\ & \nearrow \tilde{\phi} & \downarrow \\ D & \xrightarrow{\phi} & A/I \end{array}$$

commutes.

A C^* -algebra D is *semiprojective* if for any C^* -algebra A , any increasing chain of ideals I_n in A and for every $*$ -homomorphism $\phi : D \rightarrow A/I$, where $I = \overline{\bigcup I_n}$, there exist n and $*$ -homomorphism $\tilde{\phi} : D \rightarrow A/I_n$ such that the diagram

$$\begin{array}{ccc} & & A/I_n \\ & \nearrow \tilde{\phi} & \downarrow \\ D & \xrightarrow{\phi} & A/I \end{array}$$

commutes.

A C^* -algebra D is *weakly semiprojective* if for any sequence of C^* -algebras A_i and any $*$ -homomorphism $\phi : D \rightarrow \prod A_i / \bigoplus A_i$, there exists a $*$ -homomorphism $\tilde{\phi} : D \rightarrow \prod A_i$

Date: April 20, 2010.

2000 Mathematics Subject Classification. 46L05; 46L35.

Key words and phrases. Projective and semiprojective C^* -algebras, stable relation, lifting problem, M-ideal.

such that the diagram

$$\begin{array}{ccc} & & \prod A_i \\ & \nearrow \tilde{\phi} & \downarrow \\ D & \xrightarrow{\phi} & \prod A_i / \bigoplus A_i \end{array}$$

commutes.

These notions provide an algebraic setting of lifting and stability questions: a relation is liftable (stable, weakly stable) if and only if its universal C^* -algebra is projective (semiprojective, weakly semiprojective), [2], [5]. Of course it is assumed that the universal C^* -algebra of a relation exists. Note that in general a relation needn't have a universal C^* -algebra, but if it includes restrictions $\|x_i\| \leq c_i$, for all generators, then the universal C^* -algebra exists (of course if the relation has at least one representation in a Hilbert space).

Thus given a liftable or (weakly) stable relation there arises a natural question if being combined with restrictions on norms of generators it is still liftable or (weakly) stable: the positive answer would produce new examples of projective or (weakly) semiprojective C^* -algebras.

It was proved in [8] that each nilpotent is liftable and in [5], [6] it was asked a question if a relation

$$x^n = 0, \|x\| \leq 1$$

is liftable or, in other words, if it defines a projective C^* -algebra. In [13] this question was answered positively and the main tool was using the technique of M-ideals in Banach spaces.

In [4] it was proved that for each polynomial p in one variable, a relation $p(x) = 0$ is weakly stable (though the proof given in [4] easily implied also stability) but again it remained unknown if the universal C^* -algebra of a relation

$$p(x) = 0, \|x\| \leq 1$$

is semiprojective. In the present paper using approach of [13] we prove this in two "opposite" cases: when all the roots of the polynomial have multiplicity more than 1 and when all the roots have multiplicity 1.

We prove also that for polynomials with all roots multiple, the universal C^* -algebras of such relations are residually finite-dimensional (have separating families of finite-dimensional representations). Note that this in no way means that these algebras are small. For polynomials of degree more than two they all are non-exact.

Theorems 11 and 13 concern well known questions of C. Olsen. Let $T \in B(H)$, $p(x)$ be a polynomial in one variable. Olsen's questions are:

- 1) Does there exist a compact operator K such that $\|p(T + K)\| = \|p(T)\|_e$? (by $\|T\|_e$ the essential norm of T is denoted).
- 2) Can this K be chosen common for all polynomials?

The questions are still open but there are partial results ([1], [9], [10], [11], [3], [14]). In particular Olsen proved ([9]) that under the assumption that $p(T)$ is compact, such K always exists. In this paper we prove that under the same assumption K can be chosen common for polynomials $p(x)$ and x in the case when either all roots of the polynomial have multiplicity more than 1 or when they all have multiplicity 1. In other words we prove (Theorem 11) that for such polynomials if $p(T)$ is compact then there exists a compact operator K such that

$$p(T + K) = 0, \quad \|T + K\| = \|T\|_e.$$

1. PRELIMINARY RESULTS AND CONSTRUCTIONS

Below we use the following notation.

By $M(I)$ we denote the multiplier C^* -algebra of a C^* -algebra I .

For a closed ideal I of a C^* -algebra A , we denote by $a \rightarrow \dot{a}$ the standard epimorphism from A to A/I . We say that a is a lift of b if $\dot{a} = b$.

Given ideals $I_1 \subseteq I_2 \subseteq \dots \subseteq I \subseteq A$ with $I = \overline{\bigcup I_n}$ we use the notation

$$q_n : A \rightarrow A/I_n$$

and

$$q_{n,\infty} : A/I_n \rightarrow A/I$$

for the induced epimorphisms.

For an arbitrary algebra A , an operator $T : A \rightarrow A$ is called an elementary operator if it is of the form $T = \sum_{i=1}^N L_{a_i} R_{b_i}$, where $a_i, b_i \in A^+$. By L_{a_i}, R_{b_i} we denote operators of left and right multiplication by a_i and b_i respectively.

A subspace Y of a Banach space X is called proximal if for any $x \in X$ there is $y \in Y$ such that $\|x - y\| = d(x, Y)$, or, equivalently, if for any $z \in X/Y$ there is a lift x of z such that $\|z\| = \|x\|$.

Below for any C^* -algebra A we denote the set of its self-adjoint elements by A_{sa} .

Theorem 1. ([13]) *Let A be a C^* -algebra, T an elementary operator on A . Then, for each ideal I of A , $\overline{TI}$ is proximal in $\overline{TA}$ and $\overline{TI_{sa}}$ is proximal in $\overline{TA_{sa}}$.*

Remark 2. In [13] it was not considered the case of $\overline{TI_{sa}}$ but the proof is the same, if one considers C^* -algebras as Banach spaces over $\mathbb{R}$.

Theorem 1 implies a corollary which was contained in nondirect form in the proof of [Theorem 5, [13]]:

Corollary 3. *Let A be a C^* -algebra, I its ideal, T an elementary operator on A , $b \in A/I$. If, for some lift a_0 of b , an element Ta_0 is also a lift of b , then the set $\{Ta \mid \dot{a} = b\}$ contains an element of norm equal to $\|b\|$.*

Proof. Let $E = \{Ta \mid \dot{a} = b\}$. Then

$$E = \{Ta_0 + Ti \mid i \in I\}, \quad \overline{E} = \{Ta_0 + x \mid x \in \overline{TI}\}.$$

All elements of $\overline{E}$ are lifts of b . Indeed let $\dot{a} = b$, then $a = a_0 + i$, for some $i \in I$. Since T is elementary, it preserves ideals, whence $(Ti)^\cdot = 0$ and $(Ta)^\cdot = (Ta_0)^\cdot = b$. This easily implies that $\overline{E}$ also consists of lifts of b .

Let $b_i, c_i, i = 1, \dots, N$, be elements of A^+ such that $A = \sum_{i=1}^N L_{b_i} R_{c_i}$ and let B be the C^* -algebra generated by $b_i, i = 1, \dots, N$, and I . Let us choose a quasicentral approximate unit u_λ for the ideal I of B . Then, for each $a \in E$, we have $T((1 - u_\lambda)a) \in E$ and

$$\lim_\lambda \|T((1 - u_\lambda)a)\| = \lim_\lambda \|(1 - u_\lambda)Ta\| = \|(Ta)^\cdot\| = \|b\|$$

(we used here a well known fact [7], Section 3.1.3, that if e_λ is any approximate unit then

$$\lim_\lambda \|x(1 - e_\lambda)\| = \|\dot{x}\|,$$

for any $x \in A$). Hence $\inf_{y \in \overline{TI}} \|Ta_0 + y\| \leq \|b\|$. Since the norm of any lift of b is not less than $\|b\|$ we get $\inf_{y \in \overline{TI}} \|Ta_0 + y\| = \|b\|$.

Thus $\text{dist}(Ta_0, \overline{TI}) = \inf_{y \in \overline{TI}} \|Ta_0 + y\| = \|b\|$ and, by Theorem 1, there is $y_0 \in \overline{TI}$ such that $\|Ta_0 + y_0\| = \|b\|$. Since $Ta_0 + y_0 \in \overline{E}$, we are done. $\square$

Lemma 4. $\overline{\dot{a}(A/I)\dot{a}} = \overline{aAa/aIa}$.

Proof. Define a map $f : aAa \rightarrow \dot{a}(A/I)\dot{a}$ by $f(aba) = \dot{a}\dot{b}\dot{a}$ and extend it by continuity to $\tilde{f} : \overline{aAa} \rightarrow \dot{a}(A/I)\dot{a}$. Since $\tilde{f}|_{\overline{aIa}} = 0$ there is a well-defined map $F : \overline{aAa}/\overline{aIa} \rightarrow \dot{a}(A/I)\dot{a}$. This is surjection because its image is closed (as the image of any $*$ -homomorphism between C^* -algebras) and contains $\dot{a}(A/I)\dot{a}$.

Suppose that $x \in \overline{aAa}/\overline{aIa}$ and $F(x) = 0$. Let $X \in \overline{aAa}$ be any preimage of x . Then there are elements $b_n \in A$ such that $X = \lim ab_na$. Then $\lim \dot{a}b_n\dot{a} = 0$ and there are $i_n \in I$ such that $ab_na - i_n \rightarrow 0$. Hence $X \in I$. Let $\{u_\lambda\}$ be a quasicentral approximate unit for $I \subseteq A$. Then for any ϵ , there is n such that $\|X - ab_na\| \leq \epsilon$ and there is λ such that $\|X - Xu_\lambda\| \leq \epsilon$, $\|ab_nau_\lambda - ab_nu_\lambda a\| \leq \epsilon$. Then $\|X - ab_nu_\lambda a\| \leq 3\epsilon$, $\text{dist}(X, aIa) \leq 3\epsilon$ whence $X \in \overline{aIa}$. $\square$

A C^* -algebra A is called SAW* ([12]) if for any $a, b \in A_{sa}$, $ab = 0$, there is $0 \leq z \leq 1$, $z \in A^+$, such that $az = a$, $zb = 0$.

Lemma 5. *If A is SAW*, $p \in A$ is a projection, then pAp is SAW*.*

Proof. Suppose $a, b \in (pAp)_{sa}$, $ab = 0$. Since A is SAW* there is $0 \leq z \leq 1$, $z \in A^+$, such that $az = a$, $zb = 0$. Clearly $0 \leq pzp \leq 1$, $pzp \in (pAp)^+$, $apzp = a$, $pzpb = 0$. $\square$

Below we write $d \ll e$ for selfadjoint elements satisfying $de = ed = d$.

Theorem 6. ([8]) *Let A be a C^* -algebra, I — its ideal, such that the quotient A/I is SAW*. Let $b \in A/I$ be a nilpotent of order n , $n \geq 2$. Then there are elements*

$$0 \leq d_{n-1} \ll e_{n-1} \ll d_{n-2} \ll e_{n-2} \ll \dots \ll e_2 \ll d_1 \ll e_1 \leq e_0 = 1 \quad (1)$$

and their lifts

$$0 \leq D_{n-1} \ll E_{n-1} \ll D_{n-2} \ll E_{n-2} \ll \dots \ll E_2 \ll D_1 \ll E_1 \leq E_0 = 1 \quad (2)$$

such that

- (i) $b = \sum_{j=1}^{n-1} (e_{j-1} - e_j)bd_j$,
- (ii) for any lift B of b , $\sum_{j=1}^{n-1} (E_{j-1} - E_j)BD_j$ is a nilpotent lift of b of order n .

Now let an element b of a C^* -algebra be algebraic with the minimal polynomial $p(t) = (t - \lambda_1)^{k_1} \dots (t - \lambda_n)^{k_n}$. In [4] a family of n pairwise orthogonal projections was associated with b in the following way. Let e_i be the spectral idempotents for b and $s = \sum e_i^* e_i$. One can prove that $e_i(b - \lambda_i)^{k_i} e_i = 0$ for each i , s is invertible and

$$p_i = s^{1/2} e_i s^{-1/2} \quad (3)$$

are projections with sum 1. Let $c = s^{1/2} b s^{-1/2}$. Then

$$p_i(c - \lambda_i)^{k_i} p_i = 0, \quad (4)$$

$$\sum_{i=1}^n p_i c p_i = c. \quad (5)$$

Theorem 7. ([4]) *The following are equivalent:*

- (i) $b \in A/I$ has a lift $\tilde{b} \in A$ with $p(\tilde{b}) = 0$;
- (ii) the family $\{p_i\}$ can be lifted to a family of projections in A with sum 1.

Since in what follows we use ideas from [4], we now sketch arguments from [4] of how (ii) implies (i). Let a family of projections $\{Q_i\}$ with $\sum Q_i = 1$ be a lift of $\{p_i\}$. Since $p_i(c - \lambda_i)p_i$ is a nilpotent of order k_i in a C^* -algebra $p_i A / I p_i = Q_i A Q_i / Q_i I Q_i$, by [8] it has a nilpotent lift $t_i \in Q_i A Q_i$ of order k_i . Now one can check that $\sum t_i + \lambda_i Q_i$ is a lift of c and $p(\sum t_i + \lambda_i Q_i) = 0$. As a lift of b we can take $S^{-1/2}(\sum t_i + \lambda_i Q_i)S^{1/2}$, where S

is any invertible positive lift of s (to find such S we can take any selfadjoint lift of s and then use functional calculus).

2. MAIN RESULTS

Lemma 8. *Let $p(t) = (t - \lambda_1)^{k_1} \dots (t - \lambda_n)^{k_n}$ be a polynomial such that $k_i \geq 2$, $i = 1, \dots, n$, A be a C^* -algebra, I its ideal such that A/I is SAW*. Suppose that $b \in A/I$, $p(b) = 0$, $\|b\| = 1$, and a family of projections $\{p_i\}$ associated with b by (3) can be lifted to a family of projections in A with sum 1. Then there is a lift $\tilde{b} \in A$ of b such that $p(\tilde{b}) = 0$, $\|\tilde{b}\| = 1$.*

Proof. Let s, c be as above and projections $\{Q_i\}$ in A be lifts of $\{p_i\}$ with sum 1. By (4), $p_i(c - \lambda_i)p_i$ is a nilpotent of order k_i in the C^* -algebra $p_i(A/I)p_i$. By Lemma 4, $p_i(A/I)p_i = Q_i A Q_i / Q_i I Q_i$. It follows from Lemma 5 and Theorem 6 that there are $e_j^{(i)}, d_j^{(i)} \in p_i(A/I)p_i$ and their lifts $E_j^{(i)}, D_j^{(i)} \in Q_i A Q_i$ satisfying (1) and (2) such that

$$p_i(c - \lambda_i)p_i = \sum_{j=1}^{k_i-1} (e_{j-1}^{(i)} - e_j^{(i)})(c - \lambda_i)d_j^{(i)}$$

and, for any lift C of c ,

$$\sum_{j=1}^{k_i-1} (E_{j-1}^{(i)} - E_j^{(i)})(C - \lambda_i)D_j^{(i)}$$

is a nilpotent lift of $p_i(c - \lambda_i)p_i$ of order k_i . Let $q : A \rightarrow A/I$ be the canonical surjection. We have

$$\begin{aligned} q \left(\sum_{i=1}^n \sum_{j=1}^{k_i-1} \left((E_{j-1}^{(i)} - E_j^{(i)})(C - \lambda_i)D_j^{(i)} + \lambda_i Q_i \right) \right) = \\ \sum_{i=1}^n \sum_{j=1}^{k_i-1} \left((e_{j-1}^{(i)} - e_j^{(i)})(c - \lambda_i)d_j^{(i)} + \lambda_i p_i \right) = \sum_{i=1}^n p_i(c - \lambda_i)p_i + \lambda_i p_i = \sum_{i=1}^n p_i c p_i = c. \end{aligned}$$

Denote $\sum_{j=1}^{k_i-1} (E_{j-1}^{(i)} - E_j^{(i)})(C - \lambda_i)D_j^{(i)}$ by X_i . Since $X_i^{k_i} = 0$, $X_i Q_j = Q_j X_i = 0$ if $i \neq j$, and $\sum Q_i = 1$ we have

$$\begin{aligned} p \left(\sum_{i=1}^n \sum_{j=1}^{k_i-1} \left((E_{j-1}^{(i)} - E_j^{(i)})(C - \lambda_i)D_j^{(i)} + \lambda_i Q_i \right) \right) = \\ \prod_{l=1}^n \left(\sum_{i=1}^n (X_i + \lambda_i Q_i) - \lambda_l \right)^{k_l} = \prod_{l=1}^n \left(\sum_{i=1}^n (X_i + (\lambda_i - \lambda_l) Q_i) \right)^{k_l} = \\ \prod_{l=1}^n \left(X_l + \sum_{i \neq l} (X_i + \lambda_i - \lambda_l) Q_i \right)^{k_l} = \prod_{l=1}^n \left(\sum_{i \neq l} (X_i + \lambda_i - \lambda_l) Q_i \right)^{k_l} = 0. \end{aligned}$$

It follows from (2) that $\sum_{j=1}^{k_i-1} (E_{j-1}^{(i)} - E_j^{(i)})D_j^{(i)} = Q_i$ whence

$$\sum_{i=1}^n \sum_{j=1}^{k_i-1} \left((E_{j-1}^{(i)} - E_j^{(i)})(C - \lambda_i)D_j^{(i)} + \lambda_i Q_i \right) = \sum_{i=1}^n \sum_{j=1}^{k_i-1} (E_{j-1}^{(i)} - E_j^{(i)})C D_j^{(i)}.$$

Let $S \in A$ be any invertible positive lift of s . Since $b = s^{-1/2}cs^{1/2}$ the set

$$\mathcal{E} = \left\{ \sum_{i=1}^n \sum_{j=1}^{k_i-1} S^{-1/2}(E_{j-1}^{(i)} - E_j^{(i)})S^{1/2}XS^{-1/2}D_j^{(i)}S^{1/2} \mid X \in A, q(X) = b \right\}$$

consists of lifts of b and we have $p(X) = 0$, for any $X \in \mathcal{E}$. Then $\overline{\mathcal{E}}$ also consists of lifts of b and $p(X) = 0$, for any $X \in \overline{\mathcal{E}}$.

Define an elementary operator $T : A \rightarrow A$ by

$$T(X) = \sum_{i=1}^n \sum_{j=1}^{k_i-1} S^{-1/2}(E_{j-1}^{(i)} - E_j^{(i)})S^{1/2}XS^{-1/2}D_j^{(i)}S^{1/2}.$$

Then $\mathcal{E} = \{TX \mid X \in A, q(X) = b\}$. By Corollary 3, in $\overline{\mathcal{E}}$ there is an element of norm 1. $\square$

Theorem 9. *Let $p(t) = (t - \lambda_1)^{k_1} \dots (t - \lambda_n)^{k_n}$ be a polynomial such that $k_i \geq 2$, $i = 1, \dots, n$. The universal C^* -algebra of a relation $p(b) = 0, \|b\| \leq 1$ is semiprojective.*

Proof. By Theorem 14.1.7 of [5], it suffices to prove that if $b \in M(I)/I$, where $I = \overline{\bigcup I_n}$, $I_1 \subseteq I_2 \subseteq \dots$, $p(b) = 0$, $\|b\| \leq 1$, then there exists N and $\tilde{b} \in M(I)/I_N$ such that $q_{n,\infty}(\tilde{b}) = b$, $p(\tilde{b}) = 0$, $\|\tilde{b}\| \leq 1$. Here by $q_{n,\infty}$ we denote the canonical surjection from $M(I)/I_N$ onto $M(I)/I$. Clearly we can assume $\|b\| = 1$ and look for $\tilde{b}$ with $\|\tilde{b}\| = 1$.

By [2], there exists N such that $\{p_i\}$ can be lifted to a family $\{Q_i\}$ of projections in $M(I)/I_N$ with sum 1. Denote $M(I)/I$ by A , I/I_N — by J . Then $A/J \cong M(I)/I$. By Kasparov's Technical Theorem, any corona-algebra is SAW*, so the assertion follows now from Lemma 8. $\square$

Recall that a C^* -algebra is residually finite-dimensional (RFD) if the intersection of kernels of its finite-dimensional representations is zero.

Theorem 10. *Let $p(t) = (t - \lambda_1)^{k_1} \dots (t - \lambda_n)^{k_n}$ be a polynomial such that $k_i \geq 2$, $i = 1, \dots, n$. The universal C^* -algebra of a relation $p(b) = 0, \|b\| \leq 1$ is residually finite-dimensional.*

Proof. Let H be $l^2(\mathbb{N})$. We will identify M_n with $B(l^2\{1, \dots, n\}) \subseteq B(H)$. Let $\mathcal{A} \subseteq \prod M_n$ be a C^* -algebra of all sequences $\{x_n\}$ such that $*$ -strong limit x_n exists and let $\mathcal{I}$ be an ideal of all sequences $\{x_n\}$ such that $*$ -strong limit x_n is zero. Then a map $i : \mathcal{A}/\mathcal{I} \rightarrow B(H)$ sending each sequence $\{x_n\}$ to its $*$ -strong limit is an isomorphism. The algebra $B(H)$ is SAW*. Indeed if $T, S \in B(H)$, $TS = 0$ then for Z equal to the projection on $(\text{Ran } S)^\perp$ one has $TZ = T$, $ZS = 0$. Thus $\mathcal{A}/\mathcal{I}$ is SAW*. Let $p_1, \dots, p_n$ be a family of projections in $\mathcal{A}/\mathcal{I}$ with sum 1. Let H_j be a range of projection $i(p_j)$, $\{e_i^{(j)}\}_{i \in \mathbb{N}}$ be an orthonormal basis in H_j . Then $e_1^{(1)}, \dots, e_1^{(n)}, e_2^{(1)}, \dots, e_2^{(n)}, \dots$ is an orthonormal basis in H . Let R_k be a projection on the first k basis vectors. Then sequences $\{R_k P_j R_k\}_{k \in \mathbb{N}} \in \mathcal{A}$ are projections with sum 1 and are lifts of P_j . By Lemma 8 each $*$ -homomorphism from the universal C^* -algebra of a relation $p(b) = 0, \|b\| \leq 1$ to $\mathcal{A}/\mathcal{I}$ lifts to an $*$ -homomorphism to $\mathcal{A}$. Let π be the universal representation of the universal C^* -algebra of a relation $p(b) = 0, \|b\| \leq 1$. Then any lift of $i^{-1} \circ \pi$ has to be injective and hence gives a separating family of finite-dimensional representations. $\square$

Theorem 11. *Let $T \in B(H)$ and $p(t) = (t - \lambda_1)^{k_1} \dots (t - \lambda_n)^{k_n}$ be a polynomial such that $k_i \geq 2$, $i = 1, \dots, n$. Suppose that $p(T)$ is compact. Then there is a compact operator K such that $p(T + K) = 0$, $\|T + K\| = \|T\|_e$.*

Proof. Let b be the image of T in Calkin algebra $B(H)/K(H)$, $C = \|T\|_e$. Then $p(b) = 0$, $\|b\| = C$. The assertion follows now from Lemma 8, Kasparov's Technical Theorem and the well-known fact that orthogonal projections (with sum 1) in Calkin algebra can be lifted to orthogonal projections (with sum 1) in $B(H)$. $\square$

Now we consider the case of a polynomial whose roots are real and have multiplicity one.

Theorem 12. *Let $p(t) = (t - t_1) \dots (t - t_n)$ be a polynomial such that $t_i \in \mathbb{R}$, $t_i \neq t_j$ when $i \neq j$. Then the universal C*-algebra of the relations $p(x) = 0$, $\|x\| \leq C$ is semiprojective.*

Proof. Let A be a C*-algebra, $\{I_n\}$ an increasing chain of ideals of A and $I = \overline{\bigcup I_n}$. By $q_N : A \rightarrow A/I_N$ we will denote the canonical surjection and for an element a in any A/I_N , $N \in \mathbb{N}$, by $\dot{a}$ we denote its image in A/I .

Suppose $x \in A/I$, $p(x) = 0$, $\|x\| \leq C$. We have to show that x has a lift X in A/I_N , for sufficiently large N , such that $p(X) = 0$, $\|X\| \leq C$.

Let s and p_i be as in (3) and let $y = \sum_{i=1}^n t_i p_i$. By (4) and (5), $x = s^{-1/2} y s^{1/2}$. Fix any invertible lift S of s . Define $T : A \rightarrow A$ by

$$T(a) = S^{-1/2} a S^{1/2},$$

$a \in A$. Let Y_0 be any selfadjoint lift of y . We have $\|(T Y_0)\dot{}\| = \|s^{-1/2} y s^{1/2}\| = \|x\| \leq C$. Let

$$E = \{T(Y_0) + T(i) \mid i \in I_{sa}\}.$$

Let $\{e_\lambda\}$ be a quasicentral approximate unit for $I \triangleleft A$ and

$$i_\lambda = 1 - (1 - e_\lambda)^{1/2}.$$

Then $\{i_\lambda\}$ is also a quasicentral approximate unit and we have

$$\begin{aligned} \text{dist}(T(Y_0), T I_{sa}) &= \inf_{e \in E} \|e\| \leq \limsup \|T(Y_0) - T(i_\lambda Y_0 + Y_0 i_\lambda - i_\lambda Y_0 i_\lambda)\| = \\ &= \limsup \|T((1 - i_\lambda) Y_0 (1 - i_\lambda))\| = \limsup \|(T(Y_0))(1 - i_\lambda)^2\| = \\ &= \limsup \|(T(Y_0))(1 - e_\lambda)\| = \|T(Y_0)\dot{}\| \leq C. \end{aligned}$$

Since S is invertible, T is also invertible and hence $T I_{sa}$ is a closed subspace of $T A_{sa}$. By Theorem 1, there is $i_0 \in I_{sa}$ such that $\text{dist}(T Y_0, T I_{sa}) = \|T Y_0 - T i_0\|$. Let

$$Y = Y_0 - i_0.$$

Thus we found a self-adjoint lift Y of y such that

$$\|S^{-1/2} Y S^{1/2}\| \leq C.$$

Let O_i , $i = 1, \dots, n$, be neighborhoods of $t_1, \dots, t_n$ such that $O_i \cap O_j = \emptyset$, when $i \neq j$. There is $\delta > 0$ such that if $|p(t)| \leq \delta$ then $t \in \bigcup O_i$. Since $p(y) = 0$, by [Prop. 2.14, [2]] for sufficiently large N , $\|p(q_N(Y))\| \leq \delta$ and hence

$$\sigma(q_N(Y)) \subseteq \bigcup_i O_i.$$

Let D_C be the disk of radius C . Let f be a continuous function on D_C such that

- $f(t_i) = t_i$ when $t \in O_i$;
- $|f(t)| \leq C$, for any $t \in D_C$.

Let $X = q_N(S^{-1/2}f(Y)S^{1/2}) = q_N(S^{-1/2})f(q_N(Y))q_N(S^{1/2})$. Then $\dot{X} = x$. Since $f(q_N(Y))$ is normal and $\sigma(f(q_N(Y))) = \{t_1, \dots, t_n\}$, we have

$$(X - t_1) \dots (X - t_n) = q_N(S^{-1/2})(f(q_N(Y)) - t_1) \dots (f(q_N(Y)) - t_n)q_N(S^{1/2}) = 0.$$

Let p_k be polynomials that converge to f uniformly on D_C such that

$$\sup_{t \in D_C} |p_n(t)| \leq C.$$

Then

$$\begin{aligned} \|X\| &= \|q_N(S^{-1/2}f(Y)S^{1/2})\| = \lim_{k \rightarrow \infty} \|q_N(S^{-1/2})p_k(q_N(Y))q_N(S^{1/2})\| = \\ &= \lim_{k \rightarrow \infty} \|p_k(q_N(S^{-1/2}YS^{1/2}))\| \end{aligned}$$

and by the von Neumann inequality

$$\|X\| \leq \lim_{k \rightarrow \infty} \sup_{|t| \leq \|q_N(S^{-1/2}YS^{1/2})\|} |p_k(t)| \leq \lim_{k \rightarrow \infty} \sup_{t \in D_C} |p_k(t)| \leq C.$$

□

Theorem 13. *Let $p(t) = (t - t_1) \dots (t - t_n)$ be a polynomial such that $t_i \in \mathbb{R}$, $t_i \neq t_j$ when $i \neq j$ and let $T \in B(H)$. Suppose that $p(T)$ is compact. Then there is a compact operator K such that $p(T + K) = 0$, $\|T + K\| = \|T\|_e$.*

Proof. Let x be the image of T in Calkin algebra. Let $C = \|T\|_e$. It is sufficient to find a pre-image $X \in B(H)$ of x such that $p(X) = 0$ and $\|X\| = C$. As in the proof of Theorem 12 we define s , y and S and find a selfadjoint pre-image Y of y such that

$$\|S^{-1/2}YS^{1/2}\| \leq C.$$

Since $p(Y)$ is compact, its spectrum is either finite or a sequence vanishing at 0. Hence the spectrum of Y is either finite or countable with accumulation points t_i , $i = 1, \dots, n$. In either case one can find neighborhoods $O_1, \dots, O_n$ of $t_1, \dots, t_n$ respectively, such that $O_i \cap O_j = \emptyset$, when $i \neq j$, and $\sigma(Y) \subset \bigcup O_i$. Now it remains to repeat the arguments from the proof of Theorem 12. □

REFERENCES

- [1] C.A. Akemann and G.K. Pedersen, Ideal perturbations of elements in C*-algebras, Math. Scand. , 41 (1977), 117-139.
- [2] B. Blackadar, Shape theory for C*-algebras, Math. Scand. , 56 (1985), 249 - 275.
- [3] C. K. Chui, D. A. Legg, P. W. Smith, and J. D. Ward, On a question of Olsen concerning compact perturbations of operators, Michigan Math. J. 24, 1 (1977), 119-127.
- [4] D. Hadwin, Lifting algebraic elements in C*-algebras, J. Funct. Anal., 127 (1995), 431-437.
- [5] T. A. Loring, Lifting solutions to perturbing problems in C*-algebras, volume 8 of Fields Institute Monographs. American Mathematical Society, Providence, RI, 1997.
- [6] T. A. Loring, G. K. Pedersen, Smoothing techniques in C*-algebra theory, J. Operator Theory 37 (1997), 3-21.
- [7] G. J. Murphy, C*-algebras and operator theory, Academic Press, New York, 1990.
- [8] C. L. Olsen and G. K. Pedersen, Corona-algebras and their applications to lifting problems, Math. Scand. 64 (1989), 63-86.
- [9] C. L. Olsen, A structure theorem for polynomially compact operators, Amer. J. Math. 93 (1971), 686-698.
- [10] C. L. Olsen and J. K. Plastiras, Quasialgebraic operators, compact perturbations, and the essential norm, Michigan Math. J. 21 (1974), 385-397.
- [11] C. L. Olsen, Norms of compact perturbations of operators, Pacific J. Math., 68, 1 (1977).
- [12] G. K. Pedersen, C*-algebras and their Automorphism Groups, LMS Monographs 14, Academic Press, London/New-York, 1979.

- [13] T. Shulman, Lifting of nilpotent contractions, Bull. London Math. Soc., 40, 6 (2008), 1002-1006.
- [14] R. R. Smith and J. D. Ward, A note on polynomial operator approximation, Proc. Amer. Math. Soc., 88, 3 (1983).

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF COPENHAGEN, UNIVERSITETSPARKEN
5, DK-2100 COPENHAGEN, DENMARK
E-mail address: `shulman@math.ku.dk`